1. Use of a formulation of the oil-in-water emulsion type comprising an oily material, an aqueous phase and a galactolipid material as an emulsifier, as a carrier for the preparation of a topical cream or lotion providing a prolonged local effect of an incorporated pharmaceutically or cosmetically active substance.

2. Use according to claim 1, wherein the formulation comprises 0.1-50 % by weight of oily material and 0.5-20 % by weight of emulsifier.

3. Use according to claim  $\frac{1}{2}$  or 2, wherein the formulation comprises 1-40 % by weight of oily material and 0.5-10 % by weight of emulsifier.

4. Use according to any of claims 1-3, wherein the galactolipid material consists of at least 50 % by weight of digalactosyldiacylglycerols and a remainder of other polar lipids, and constitutes an amount of 1.0-5.0 % by weight of the formulation.

5. Use according to any of claims 1-4, wherein the galactolipid material consists of 50-70 % by weight of digalactosyldiacylglycerols and 30-50 % by weight of other polar lipids.

6. Use according to any of claims 1-3, wherein the galactolipid material is a fractionated oat oil which consists of at least 15% by weight of digalactosyldiacylglycerols and a remainder of other polar and non-polar lipids, and constitutes an amount of 2.0-10% by weight of the formulation.

7. Use according to any of claims 1-3 and 6, wherein the galactolipid material is a fractionated oat oil which contains 40-60 % by weight polar lipids and a remainder of non-polar lipids.

2 0 -06- 2000

8. Use according to any of claims 1-7, of a cream base, comprising, in % by weight

Oily material

10.0-30.0 %

Galactolipid emulsifier

0.5-5 %

Thickener

2.0-10.0 %

Preservative

0.1-1.0 %

Water

ad 100 %

Suis

m

Con the first time con

9. Use according to any of claims 1-8 for the preparation of a topical cream or lotion, incorporating a moisturiser, especially glycerol, as the active substance.

10. Use according to any of claims 1-9 for the preparation of a medicament for prophylaxis or treatment of atopic dermatitis.

11. Use according to any of claims 1-8 for the preparation of a topical cream or lotion, incorporating a corticosteroid as the active substance, for treatment of skin inflammation.

12. Use according to any of claims 1-8, for the preparation of a topical anti-psoriatic cream or lotion, incorporating 13-hydroxy-linoleic acid as the active substance.

ACT /